

USGS-NPS Vegetation Mapping Program
Effigy Mounds National Monument

***Populus deltoides* - *Salix nigra* Forest**

COMMON NAME	Eastern Cottonwood - Black Willow Forest
SYNONYM	Midwestern Cottonwood - Black Willow Forest
PHYSIOGNOMIC CLASS	Forest (I)
PHYSIOGNOMIC SUBCLASS	Deciduous forest (I.B)
PHYSIOGNOMIC GROUP	Cold-deciduous forest (I.B.2)
PHYSIOGNOMIC SUBGROUP	Natural/Semi-natural Cold-deciduous forest (I.B.2.N)
FORMATION	Temporarily flooded cold-deciduous forest (I.B.2.N.d)
ALLIANCE	POPULUS DELTOIDES TEMPORARILY FLOODED FOREST ALLIANCE

CLASSIFICATION CONFIDENCE LEVEL 2

USFWS WETLAND SYSTEM Temporarily flooded cold-deciduous forest

CONCEPT SUMMARY

Globally

This cottonwood - black willow forest is characteristic of the fronts and banks of most major rivers and streams throughout the Central Forest Region, extending into the northern forest particularly within the Mississippi, Ohio, and Missouri River systems. It develops on bare, moist soil on recently formed sand bars, front-land ridges, and well-drained flats, along with *Salix interior*, *Eragrostis hypnoides*, *Leptochloa panicea* ssp. *brachiata* (= *Leptochloa filiformis*), *Lipocarpha micrantha* (= *Hemicarpha micrantha*), *Rumex maritimus*, *Potentilla paradoxa*, and *Bidens* spp. This natural community can also be found on abandoned fields and well-drained ridges in the first bottoms. Soils are formed in alluvium, are deep, medium-textured, and with adequate or excessive moisture available for vegetation during the growing season. The tree canopy is tall (to 30 m) and dominated by *Populus deltoides* and *Salix nigra*, although *Fraxinus pennsylvanica*, *Acer saccharinum*, *Acer negundo*, *Platanus occidentalis*, and *Ulmus americana* are also commonly encountered. Tree diversity is limited due to the dynamics of flooding and resultant deposition and scouring of sediments. The subcanopy is almost exclusively *Salix nigra*. The shrub layer is conspicuously absent in many parts of the range. Herbaceous growth can be thick and lush but is often patchy and sparse due to frequent inundation. Species most often encountered in the ground layer include *Carex* spp., *Leersia oryzoides*, *Bidens* spp., and Asteraceae spp.

RANGE

Effigy Mounds National Monument

Globally

This cottonwood - black willow forest is characteristic of the fronts and banks of most major rivers and streams throughout the Central Forest Region, extending into the northern forest particularly within the Mississippi, Ohio, and Missouri River systems, extending from Ohio west to Minnesota, southward to Oklahoma, and east to Kentucky. This community once occupied vast tracts of land along river fronts and floodplain depressions. Land clearing, ditching and draining for conversion to cropland, and logging have eliminated much of the presettlement stands of this natural community.

ENVIRONMENTAL DESCRIPTION

Effigy Mounds National Monument

Globally

This community is quick to colonize newly deposited substrates adjacent to rivers, lakes, streams, and in frequently flooded, low, wet depressions in floodplains. Dynamic substrate availability caused by frequent flooding encourages the establishment and maintenance of this community type.

MOST ABUNDANT SPECIES

Effigy Mounds National Monument

Stratum

Species

Globally

Stratum

Species

TREE CANOPY *Populus deltoides*, *Salix nigra*

TREE SUB-CANOPY *Salix nigra*

GRAMINOID *Carex typhina*, *Leersia oryzoides*

FORB *Bidens aristosa*, *Spermacoce glabra*, *Symphyotrichum lanceolatum* var *lanceolatum*,
Symphyotrichum lateriflorum var *lateriflorum*

CHARACTERISTIC SPECIES

Effigy Mounds National Monument

USGS-NPS Vegetation Mapping Program

Effigy Mounds National Monument

Globally

VEGETATION DESCRIPTION

Effigy Mounds National Monument

Vegetation of this type was not characterized during this project. However, based on observations, species likely to occur include *Populus deltoides*, *Salix nigra*, *Acer saccharinum*, and *Fraxinus pennsylvanica*.

Globally

This community is dominated by broadleaf deciduous trees. Canopy closure is complete, or nearly so, with few shrubs and limited tree species found in the type. The tree canopy is tall (to 30 m) and dominated by *Populus deltoides* and *Salix nigra*, although *Fraxinus pennsylvanica*, *Acer saccharinum*, *Acer negundo*, *Platanus occidentalis*, and *Ulmus americana* are also commonly encountered. Tree diversity is limited due to the dynamics of flooding and deposition/scouring of sediments. The subcanopy is almost exclusively *Salix nigra*. The shrub layer is conspicuously absent in many parts of the range. Herbaceous growth can be thick and lush but is often patchy and sparse due to frequent inundation. Species most often encountered in the ground layer include *Carex* spp., *Leersia oryzoides*, *Bidens* spp., and Asteraceae spp. (TNC 1995a).

Species composition is uniform throughout the range of this community. Species density is governed by the duration and depth of flooding. The more stable sites display very large cottonwood trees with lush understory and herbaceous layers. Sites frequently affected by flooding exhibit dense even-aged stands of cottonwood and willow. This forest often has considerable deposits of woody debris and high tree mortality.

OTHER NOTEWORTHY SPECIES

CONSERVATION RANK G3G4. The current range of this community is much smaller than the presettlement range due to extensive logging, ditching, draining, and land clearing for conversion to croplands. The reduced water flows and channelization of rivers decreases the frequency of natural floods necessary for the scouring and deposition of new substrates that favor cottonwood regeneration.

DATABASE CODE CEGL002018

COMMENTS

Effigy Mounds National Monument

Globally

REFERENCES

- Braun, E. L. 1950. Deciduous forests of eastern North America. Hafner Press, New York. 596 pp.
- Bruner, W. E. 1931. The vegetation of Oklahoma. Ecological Monographs 1:99–188.
- Collins, S. L., P. G. Risser, and E. L. Rice. 1981. Ordination and classification of mature bottomland forests in north central Oklahoma. Bulletin of the Torrey Botanical Club 108:152–165.
- Cowardin, L. M., V. Carter, F. C. Golet, and E. T. LaRoe. 1979. Classification of wetlands and deepwater habitats of the United States. U.S. Fish and Wildlife Service, Biological Service Program. FWS/OBS-79/31. Washington, DC. 103 pp.
- Evans, M. 1991. Kentucky ecological communities. Draft report to the Kentucky Nature Preserves Commission. 19 pp.
- Eyre, F. H., editor. 1980. Forest cover types of the United States and Canada. Society of American Foresters, Washington, DC. 148 pp.
- Galloway, L. A. 1963. The vegetation of an actively eroding canyon in Canadian County, Oklahoma. Unpublished M.S. thesis, University of Oklahoma, Norman.
- Galloway, L. A. 1964. The vegetation of an actively eroding canyon in Canadian County, Oklahoma. Proceedings of the Oklahoma Academy of Science 45:20–23.
- Hefley, H. M. 1937. Ecological studies on the Canadian River floodplain in Cleveland County, Oklahoma. Ecological Monographs 7:347–402.
- Hoagland, B. 2000. The vegetation of Oklahoma: A classification for landscape mapping and conservation planning. The Southwestern Naturalist 45(4):385–420.
- Kuchler, A. W. 1964. Potential natural vegetation of the conterminous United States. American Geographic Society Special Publication 36. New York, NY. 116 pp.
- Nelson, P. W. 1985. The terrestrial natural communities of Missouri. Missouri Natural Areas Committee, Jefferson City. 197 pp. Revised edition, 1987.
- Rogers, C. M. 1953. The vegetation of the Mesa de Maya region of Colorado, New Mexico, and Oklahoma. Lloydia 16(4):257–290.
- Society of American Foresters. 1967. Forest cover types of North America. Society of American Foresters, Washington, DC. 67 pp.
- TNC [The Nature Conservancy]. 1985. Global Vertebrate Characterization Abstract Habitats. Unpublished document. The Nature Conservancy, Arlington, VA.

USGS-NPS Vegetation Mapping Program
Effigy Mounds National Monument

- TNC [The Nature Conservancy]. 1995a. A classification and description of plant communities in southern Illinois. Report by the Southern Illinois Field Office, Ullin, IL, and the Midwest Regional Office, Minneapolis, MN.
- UNESCO [United Nations Educational, Scientific and Cultural Organization]. 1973. International classification and mapping of vegetation. Series 6, Ecology and Conservation. United Nations Educational, Scientific, and Cultural Organization. Paris. 93 pp.
- Voigt, J. W., and R. H. Mohlenbrock. 1964. Plant communities of southern Illinois. Southern Illinois University Press, Carbondale. 202 pp.
- Ware, G. H., and W. T. Penfound. 1949. The vegetation of the lower levels of the floodplain of the south Canadian River in central Oklahoma. *Ecology* 30:478–484.
- White, J., and M. Madany. 1978. Classification of natural communities in Illinois. Pages 311–405 in: Natural Areas Inventory technical report: Volume I, survey methods and results. Illinois Natural Areas Inventory, Urbana, IL.
- Zanoni, T. A., P. G. Risser, and I. H. Butler. 1979. Natural areas for Oklahoma. Oklahoma Natural Heritage Program, Norman. 72 pp.